

## Relationship of body mass index with efficacy of exenatide twice daily added to insulin glargine in patients with type 2 diabetes

This *post hoc* analysis assessed the evidence behind common reimbursement practices by evaluating the relationship of body mass index (BMI) ranges (<30, 30–35 and >35 kg/m<sup>2</sup>) with treatment effects of exenatide twice daily among patients with type 2 diabetes. Patients received exenatide twice daily added to insulin glargine in two 30-week studies (exenatide twice daily vs insulin lispro, n = 627; exenatide twice daily vs placebo, n = 259). No association of baseline BMI with changes in efficacy variables was observed. Glycated haemoglobin (HbA1c) reductions were significant ( $p < 0.0001$ ) and similar across BMI range groups in the lispro-comparator study and greater for exenatide versus placebo in the placebo-controlled study. Significant weight loss occurred with exenatide across BMI range groups ( $p < 0.0001$ ), while weight increased with both comparators. Achievement of HbA1c <7.0% (<53 mmol/mol) without weight gain was greater for exenatide versus comparators. Systolic blood pressure decreased across BMI range groups with exenatide in the lispro-comparator study ( $p < 0.0001$ ); changes in lipids were not clinically meaningful. Minor hypoglycaemia was less frequent for exenatide versus insulin lispro. These findings suggest that BMI alone should not limit clinical decision-making or patient access to medication.

**Keywords:** body mass index, exenatide twice daily, type 2 diabetes

Date submitted 6 January 2016; date of first decision 25 January 2016; date of final acceptance 22 March 2016

### Introduction

Type 2 diabetes (T2D) is a progressive disease requiring stepwise treatment intensification to maintain glycaemic control [1]. Recently, we reported that adding the short-acting glucagon-like peptide-1 receptor agonist (GLP-1RA) exenatide twice daily to basal insulin improved glycaemic control similarly to the addition of mealtime (three times daily) insulin lispro to basal insulin, but with less non-nocturnal hypoglycaemia, weight loss, reduced systolic blood pressure (SBP) and improved quality of life [2]. Additional studies have reported beneficial effects of the combination of a GLP-1RA and insulin [3,4]. Accordingly, recent international guidelines have included the addition of a GLP-1RA to background basal insulin as an option for patients requiring combination injectable treatments [1]. However, several countries restrict reimbursement of GLP-1RAs to only patients with a body mass index (BMI)  $\geq 35$  kg/m<sup>2</sup> [5,6].

In the present *post hoc* analysis, we assessed the evidence behind common reimbursement practices by evaluating the relationship of standard BMI ranges with the effects of exenatide twice daily among patients with T2D treated with insulin glargine in two 30-week studies.

### Materials and Methods

#### Study Design and Patients

This *post hoc* analysis included two open-label, randomized, non-inferiority studies that were published previously [2,7]. The lispro-comparator study (NCT00960661) investigated the efficacy and safety of exenatide twice daily versus insulin lispro added to background insulin glargine and metformin, after a basal insulin optimization phase in which bedtime insulin glargine was titrated based on self-monitored blood glucose. The placebo-controlled study (NCT00765817) investigated the efficacy and safety of exenatide twice daily versus placebo when added to background insulin glargine, with or without metformin and/or pioglitazone. Detailed methods of both studies have been described previously [2,7], and study procedures are shown in Figure S1, Supporting Information. The exenatide dose was 5  $\mu$ g twice daily for the first 4 weeks and 10  $\mu$ g twice daily thereafter, given before breakfast and dinner.

The protocol for each study was approved by an institutional review board, and both studies were conducted in accordance with the principles described in the Declaration of Helsinki. All patients provided written informed consent.

#### Statistical Analyses

The primary objective was to compare the changes from baseline in efficacy and safety measures with add-on exenatide twice daily versus insulin lispro or placebo for three groups with standard ranges of baseline BMI (<30, 30–35 and >35 kg/m<sup>2</sup>) [8,9] in the intention-to-treat (ITT) population,

Correspondence to: B. H. R. Wolffenbuttel, Department of Endocrinology, University of Groningen, University Medical Center Groningen, HPC AA31, Groningen 9700 RB, The Netherlands.

E-mail: bwo@umcg.nl

[The copyright line for this article was changed on 6th June after original online publication.]

This is an open access article under the terms of the Creative Commons Attribution-NonCommercial License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited and is not used for commercial purposes.

**Table 1.** Baseline characteristics by baseline BMI range group (intention-to-treat population).

Lispro-comparator study characteristics	Exenatide twice daily			Insulin lispro		
	<30 kg/m <sup>2</sup> n = 107	30–35 kg/m <sup>2</sup> n = 120	>35 kg/m <sup>2</sup> n = 88	<30 kg/m <sup>2</sup> n = 113	30–35 kg/m <sup>2</sup> n = 111	>35 kg/m <sup>2</sup> n = 88
Male, n (%)	68 (63.6)	52 (43.3)	35 (39.8)	55 (48.7)	62 (55.9)	41 (46.6)
Age, years	61.8 (9.2)	59.0 (10.2)	59.1 (8.6)	59.0 (9.5)	60.8 (9.2)	59.0 (9.7)
Body weight, kg	77.0 (9.6)	89.5 (11.5)	106.1 (15.4)	76.3 (10.6)	90.0 (12.0)	105.0 (16.5)
BMI, kg/m <sup>2</sup>	27.3 (1.7)	32.4 (1.4)	38.7 (2.5)	27.6 (1.7)	32.1 (1.3)	38.4 (2.6)
HbA1c, mmol/mol	65 (9.5)	68 (11.1)	68 (11.6)	65 (9.1)	66 (9.6)	68 (10.1)
FPG, mmol/l	6.4 (2.0)	7.5 (2.6)	7.5 (2.3)	6.4 (2.3)	7.4 (2.5)	7.3 (2.7)
SBP, mmHg	135.6 (14.5)	136.6 (16.3)	140.0 (16.3)	130.9 (15.9)	134.9 (15.1)	137.9 (14.4)
DBP, mmHg	75.9 (9.5)	79.9 (8.5)	81.7 (10.3)	76.0 (9.3)	78.4 (8.5)	80.7 (9.2)
LDL cholesterol, mmol/l	2.48 (0.92)	2.53 (0.86)	2.58 (0.83)	2.63 (0.88)	2.51 (0.89)	2.66 (0.90)
HDL cholesterol, mmol/l	1.29 (0.38)	1.21 (0.32)	1.19 (0.27)	1.26 (0.32)	1.14 (0.32)	1.20 (0.31)
Total cholesterol, mmol/l	4.55 (1.1)	4.57 (1.0)	4.61 (0.95)	4.63 (0.99)	4.48 (1.13)	4.68 (0.99)
Triglycerides, mmol/l, median (range)	1.49 (0.50, 5.97)	1.59 (0.51, 9.13)	1.65 (0.61, 7.57)	1.45 (0.67, 5.54)	1.48 (0.57, 8.64)	1.60 (0.60, 5.73)

  

Placebo-controlled study characteristics	Exenatide twice daily			Placebo		
	<30 kg/m <sup>2</sup> n = 36	30–35 kg/m <sup>2</sup> n = 50	>35 kg/m <sup>2</sup> n = 51	<30 kg/m <sup>2</sup> n = 43	30–35 kg/m <sup>2</sup> n = 35	>35 kg/m <sup>2</sup> n = 44
Male, n (%)	19 (52.8)	32 (64.0)	19 (37.3)	32 (74.4)	22 (62.9)	24 (54.5)
Age, years	62.2 (8.7)	57.7 (9.5)	55.5 (7.5)	60.3 (12.0)	59.5 (7.7)	56.7 (9.1)
Body weight, kg	73.7 (12.8)	94.6 (11.4)	111.4 (17.1)	75.6 (12.8)	89.7 (11.3)	113.4 (16.2)
BMI, kg/m <sup>2</sup>	26.8 (2.6)	32.7 (1.4)	39.9 (3.2)	26.9 (2.6)	32.3 (1.3)	39.7 (3.9)
HbA1c, mmol/mol	67 (8.0)	66 (8.6)	69 (10.5)	69 (11.0)	70 (10.5)	70 (10.2)
FPG, mmol/l	7.2 (2.8)	7.2 (2.1)	7.5 (2.8)	6.6 (2.2)	7.8 (3.3)	8.0 (2.4)
SBP, mmHg	129.8 (15.0)	128.4 (15.5)	131.4 (17.4)	126.4 (12.3)	126.6 (13.7)	129.5 (15.1)
DBP, mmHg	71.5 (8.1)	75.5 (8.5)	79.0 (10.7)	72.7 (7.9)	74.0 (10.6)	76.2 (7.9)
LDL cholesterol, mmol/l	2.61 (0.95)	2.32 (0.93)	2.55 (0.92)	2.31 (0.69)	2.43 (0.83)	2.55 (1.05)
HDL cholesterol, mmol/l	1.20 (0.30)	1.12 (0.29)	1.13 (0.24)	1.19 (0.31)	1.13 (0.29)	1.09 (0.30)
Total cholesterol, mmol/l	4.57 (1.10)	4.22 (1.07)	4.55 (1.09)	4.12 (0.88)	4.34 (0.90)	4.43 (1.17)
Triglycerides, mmol/l, median (range)	1.40 (0.45, 5.68)	1.49 (0.59, 5.31)	1.83 (0.45, 5.57)	1.27 (0.46, 3.79)	1.50 (0.73, 4.03)	1.46 (0.80, 3.63)

Data are mean (standard deviation) unless otherwise stated. BMI, body mass index; DBP, diastolic blood pressure; FPG, fasting plasma glucose; HbA1c, glycated haemoglobin; SBP, systolic blood pressure.

defined as all randomized patients who received ≥1 dose of study drug.

Efficacy measures included glycated haemoglobin (HbA1c), fasting plasma glucose (FPG), body weight, blood pressure (BP), LDL cholesterol, HDL cholesterol, LDL cholesterol/HDL cholesterol ratio, total cholesterol, triglycerides, and achievement of HbA1c <7.0% (<53 mmol/mol) and HbA1c <7.0% with no weight gain. Changes in BP and lipids were determined for patients in each BMI range group who did not add new antihypertensive or lipid-lowering medication, respectively. Safety measures included the incidence of treatment-emergent adverse events (TEAEs) and minor and major hypoglycaemia.

Baseline demographics, TEAEs, hypoglycaemia rates and goal achievement were summarized by descriptive statistics. Changes from baseline to endpoint in efficacy variables were evaluated with the last observation carried forward method and analysed with an analysis of covariance model, in which the baseline value of the dependent variable was adjusted as a covariate. The relationship of baseline BMI to the change from baseline to endpoint in efficacy variables was examined, and the Pearson correlation coefficients (*r*) were provided.

## Results

### Study Population

A total of 627 of 637 randomized patients in the lispro-comparator study and 259 of 261 randomized patients in the placebo-controlled study received ≥1 dose of study medication after randomization and were included in this analysis. Mean baseline HbA1c ranged from 8.1 to 8.6% (65 to 69 mmol/mol; Table 1).

### Efficacy Measures

HbA1c was significantly reduced from baseline among all groups (*p* < 0.001; Table 2). FPG reductions were numerically greater for exenatide twice daily versus comparator in both studies in all BMI range groups except BMI <30 kg/m<sup>2</sup> in the placebo-controlled study. Baseline BMI was not correlated with changes from baseline in HbA1c (range of *r*-values, −0.05 to −0.15) or FPG (range of *r*-values, −0.04 to −0.13) for any treatment group in either study. Achievement of HbA1c <7.0% (<53 mmol/mol) was numerically higher for patients receiving add-on exenatide twice daily versus comparator in all groups

Table 2. Metabolic and cardiovascular characteristics in the intention-to-treat population.

	Exenatide twice daily		Insulin lispro	
	BMI <30 kg/m <sup>2</sup> n = 107	BMI 30–35 kg/m <sup>2</sup> n = 120	BMI <30 kg/m <sup>2</sup> n = 113	BMI >35 kg/m <sup>2</sup> n = 88
<b>Lispro-comparator study</b>				
Change from baseline				
HbA1c, mmol/mol	-10.7 (1.0)*	-10.9 (0.8)*	-9.6 (0.9)*	-11.0 (0.9)*
FPG, mmol/l	-0.22 (0.26)	-0.65 (0.27)*	0.67 (0.28)*	0.22 (0.32)
Body weight, kg	-2.1 (0.3)*	-2.0 (0.3)*	1.8 (0.3)*	1.5 (0.3)*
SBP, mmHg	-4.3 (1.5)*	-4.0 (1.4)*	2.1 (1.6)	1.8 (1.3)
DBP, mmHg	-0.2 (0.9)	-1.1 (0.8)	1.3 (1.0)	0.6 (0.8)
LDL cholesterol, mmol/l	-0.01 (0.07)	-0.11 (0.05)*	0.05 (0.06)	-0.01 (0.06)
HDL cholesterol, mmol/l	-0.05 (0.02)*	-0.04 (0.02)*	0.03 (0.02)	0.05 (0.02)*
LDL cholesterol/HDL cholesterol ratio, mmol/l	0.07 (0.06)	-0.04 (0.05)	0.01 (0.06)	-0.09 (0.06)
Total cholesterol, mmol/l	-0.09 (0.08)	-0.1 (0.06)	0.09 (0.08)	0.05 (0.07)
Triglycerides, mmol/l, median	-0.07	0.12	0.01	-0.05
Patients achieving HbA1c <7.0% (<53 mmol/mol) at endpoint, n (%)	56 (52.3)	42 (35.0)	46 (40.7)	51 (45.9)
Patients achieving HbA1c <7.0% (<53 mmol/mol) and no weight gain at endpoint, n (%)	51 (47.7)	31 (25.8)	16 (14.2)	21 (18.9)
<b>Placebo-controlled study</b>				
Change from baseline				
HbA1c, mmol/mol	-12.5 (2.0)*	-16.5 (1.6)*	-9.6 (1.6)*	-11.9 (1.5)*
FPG, mmol/l	-0.32 (0.65)	-1.45 (0.39)*	-0.53 (0.45)	-0.94 (0.65)
Body weight, kg	-0.4 (0.4)	-2.6 (0.6)*	1.2 (0.4)*	1.2 (0.6)*
SBP, mmHg	-3.6 (2.5)	-2.8 (2.3)	1.2 (2.0)	1.6 (3.0)
DBP, mmHg	-1.5 (1.0)	-2.1 (1.2)	3.0 (1.0)*	1.2 (1.6)
LDL cholesterol, mmol/l	0.0 (0.17)	-0.30 (0.10)*	0.14 (0.10)	0.04 (0.10)
HDL cholesterol, mmol/l	-0.05 (0.03)	0 (0.02)	-0.01 (0.09)	0 (0.03)
LDL cholesterol/HDL cholesterol ratio, mmol/l	0.11 (0.15)	-0.31 (0.11)*	0.13 (0.13)	0.05 (0.08)
Total cholesterol, mmol/l	0.03 (0.18)	-0.28 (0.11)*	0.14 (0.11)	0.08 (0.11)
Triglycerides, mmol/l, median	0.10	-0.17	-0.02	-0.02
Patients achieving HbA1c <7.0% (<53 mmol/mol) at endpoint, n (%)	16 (44.4)	33 (66.0)	11 (25.6)	10 (28.6)
Patients achieving HbA1c <7.0% (<53 mmol/mol) and no weight gain at endpoint, n (%)	10 (27.8)	25 (50.0)	5 (11.6)	8 (22.9)

Data are mean (standard error) unless otherwise stated. BMI, body mass index; BP, blood pressure; DBP, diastolic blood pressure; FPG, fasting plasma glucose; HbA1c, glycated haemoglobin; SBP, systolic blood pressure.  
\*p < 0.05.

except for the BMI 30–35 kg/m<sup>2</sup> group in the lispro-comparator study (Table 2).

Body weight was significantly reduced among patients receiving add-on exenatide twice daily across all groups except the <30 kg/m<sup>2</sup> BMI group in the placebo-controlled study ( $p < 0.05$ ; Table 2). Increases in body weight were observed for both comparators. Baseline BMI was not correlated with changes from baseline in body weight for any treatment in either study (range of  $r$ -values,  $-0.18$  to  $0.09$ ). Achievement of HbA1c  $< 7.0\%$  ( $< 53$  mmol/mol) and no weight gain was consistently higher for patients receiving exenatide twice daily versus comparator in both studies (Table 2).

When excluding patients who added on new antihypertensive medications (lispro-comparator study: exenatide twice daily,  $n = 2$ ; insulin lispro,  $n = 5$ ; placebo-controlled study: exenatide twice daily,  $n = 14$ ; placebo,  $n = 14$ ), the BP-lowering effect of add-on exenatide twice daily remained and changes in BP observed were similar to those in the overall ITT populations of the BMI range groups (Table 2). Baseline BMI was not correlated with changes from baseline in SBP (range of  $r$ -values,  $-0.03$  to  $-0.11$ ) or diastolic BP (range of  $r$ -values,  $-0.08$  to  $-0.14$ ) for any treatment in either study.

None of the patients in the lispro-comparator study added on new lipid-lowering medications after randomization. In the placebo-controlled study, 10 patients receiving exenatide twice daily and zero placebo-treated patients added on lipid-lowering medications. When excluding these patients, the changes from baseline in LDL cholesterol, HDL cholesterol and total cholesterol were similar to those in the overall ITT population of each BMI range group. Baseline BMI was not correlated with changes in LDL cholesterol, HDL cholesterol, total cholesterol, or triglycerides for any treatment in either study (range of  $r$ -values for all lipids,  $-0.14$  to  $0.24$ ).

### Safety and Tolerability

Rates of minor hypoglycaemia were numerically lower among patients receiving add-on exenatide twice daily versus add-on insulin lispro in all BMI range groups in the lispro-comparator study and versus placebo in the lowest BMI range groups in the placebo-controlled study (Table S1, Supporting Information). Major hypoglycaemia was infrequent and occurred in a smaller proportion of patients receiving exenatide twice daily versus insulin lispro in the lispro-comparator study; major hypoglycaemia was not reported in the placebo-controlled study for exenatide twice daily. The most frequent TEAEs ( $\geq 10\%$  incidence) by preferred term included diarrhoea, nausea, vomiting and nasopharyngitis in the lispro-comparator study, and nausea, diarrhoea and vomiting in the placebo-controlled study.

### Discussion

In this secondary analysis of two clinical trials in which patients with T2D were randomized to add-on exenatide twice daily versus add-on insulin lispro or placebo, while taking background insulin glargine with or without metformin and/or pioglitazone, baseline BMI was not correlated with changes from baseline in the efficacy variables evaluated. In both studies, exenatide twice daily was associated with

significant reductions in HbA1c and body weight. In the lispro-comparator study, HbA1c reductions were equivalent for exenatide twice daily and insulin lispro; however, insulin lispro was associated with significant weight gain. Thus, the decision by some European national authorities or health insurance companies to restrict GLP-1RA reimbursement to patients with BMI  $> 35$  kg/m<sup>2</sup> is not consistent with medical evidence.

Other studies of exenatide twice daily have also found similar efficacy across BMI subgroups. In *post hoc* analyses of exenatide twice daily (16 randomized controlled trials), HbA1c, body weight and SBP reductions were significant for all BMI range groups ( $p < 0.0001$ ) [10,11]. Similarly, studies with liraglutide [12] and lixisenatide added on to basal insulin [13] have reported reductions in HbA1c and body weight across BMI range groups. These effects appear to complement the guideline-recommended management of cardiovascular risk factors.

Hypoglycaemia is a concern in patients with T2D taking both basal and mealtime insulins [14,15], and exenatide twice daily added to background insulin glargine appeared to mitigate this risk. The trend of fewer minor hypoglycaemic events in patients with BMI  $< 30$  kg/m<sup>2</sup> receiving exenatide twice daily treatment versus comparators in both studies is important because low BMI (25 vs 35 kg/m<sup>2</sup>) is associated with increased non-severe hypoglycaemia risk in patients with T2D [16]. In both studies and for all BMI range groups, exenatide twice daily treatment was associated with an increased rate of gastrointestinal TEAEs versus comparator.

Limitations of the present analysis include differences in study design, the retrospective nature of the analysis, and the lack of balance across BMI range groups.

In conclusion, recent international guidelines have adopted a more favourable opinion on the combination of basal insulin and GLP-1RAs after basal insulin failure. Exenatide twice daily effectively reduced HbA1c and body weight across BMI range groups, including patients with near-normal BMI, and was well tolerated with a low risk of hypoglycaemia. Changes in efficacy variables were not significantly correlated with baseline BMI. There is evidence of clinically meaningful glycaemic and weight benefits from the addition of exenatide twice daily to insulin glargine in both patients of normal weight and patients with mild to severe obesity, and there does not appear to be a threshold below which exenatide twice daily is not effective. Limiting reimbursement to patients with obesity (BMI  $> 35$  kg/m<sup>2</sup>) may result in the undertreatment of patients with lower BMI and may increase their risk of hypoglycaemia and unwanted weight gain.

**B. H. R. Wolffenbuttel<sup>1</sup>, L. Van Gaal<sup>2</sup>, S. Durán-García<sup>3</sup>  
& J. Han<sup>4</sup>**

<sup>1</sup>Department of Endocrinology, University of Groningen, University Medical Center Groningen, Groningen, The Netherlands

<sup>2</sup>Department of Endocrinology, Diabetology, and Metabolism, Antwerp University Hospital, Antwerp, Belgium

<sup>3</sup>Unidad de Gestión de Endocrinología y Nutrición, Hospital Universitario Virgen de Valme, Seville, Spain

<sup>4</sup>Pharmapace, Inc, San Diego, CA, USA

## Acknowledgements

The studies included in the analysis were supported by the alliance of Eli Lilly and Company and Amylin Pharmaceuticals and the alliance of Bristol-Myers Squibb and AstraZeneca. Manuscript development was supported by AstraZeneca. Raewyn Poole and Robert Schupp, PharmD, CMPP, of inScience Communications, Springer Healthcare (Philadelphia, PA, USA), provided medical writing support funded by AstraZeneca.

## Conflict of Interest

B. H. R. W. has received grant support for clinical studies and also consulting fees for serving on advisory boards and has served as a speaker for AstraZeneca, Eli Lilly and Company, GlaxoSmithKline, Novo Nordisk, Pfizer and Sanofi Aventis. He has also received consulting fees from Eli Lilly and Company as a member of the 4B Study and of the DURABLE Trial Data Monitoring Committee, and grant support from the European Union (BBMRI, BioSHaRE, and MDS-Right Consortiums), JDRF and Diabetes Fonds Nederland. L. V. G. is/has been a member of the advisory boards and speakers' bureaus for AstraZeneca/BMS, Boehringer Ingelheim, Eli Lilly, Janssen, Johnson & Johnson, Merck MSD, Novo Nordisk and Sanofi (2011–2013). He also received grant support from the European Union (Hepadip & Resolve Consortium) and National Research Funds, Belgium. S. D. G. has received grant support for clinical studies from AstraZeneca, Eli Lilly and Company, GlaxoSmithKline, Intarcia, Lexicon, Novo Nordisk, Pfizer, Servier, and Takeda. J. H. has received consulting fees for statistical consulting.

B. H. R. W. and L. V. G. were investigators in the 4B Study Group, B. H. R. W., L. V. G., S. D. G. and J. H. conceived the analysis and participated in the writing of the manuscript. All authors approved the final version of the manuscript.

## Supporting Information

Additional Supporting Information may be found in the online version of this article:

[Figure S1](#). Study design.

[Table S1](#). Treatment-emergent adverse events occurring in  $\geq 10\%$  of patients in any group and hypoglycaemic events, according to baseline body mass index range group (intent-to-treat population).

## References

- Inzucchi SE, Bergenstal RM, Buse JB et al. Management of hyperglycemia in type 2 diabetes, 2015: a patient-centered approach: update to a position statement of the American Diabetes Association and the European Association for the Study of Diabetes. *Diabetes Care* 2015; **38**: 140–149.
- Diamant M, Nauck MA, Shaginian R et al. Glucagon-like peptide 1 receptor agonist or bolus insulin with optimized basal insulin in type 2 diabetes. *Diabetes Care* 2014; **37**: 2763–2773.
- Gough SC, Bode B, Woo V et al. Efficacy and safety of a fixed-ratio combination of insulin degludec and liraglutide (IDegLira) compared with its components given alone: results of a phase 3, open-label, randomised, 26-week, treat-to-target trial in insulin-naïve patients with type 2 diabetes. *Lancet Diabetes Endocrinol* 2014; **2**: 885–893.
- Rosenstock J, Fonseca VA, Gross JL et al. Advancing basal insulin replacement in type 2 diabetes inadequately controlled with insulin glargine plus oral agents: a comparison of adding albiglutide, a weekly GLP-1 receptor agonist, versus thrice-daily prandial insulin lispro. *Diabetes Care* 2014; **37**: 2317–2325.
- Divino V, DeKoven M, Hallinan S et al. Glucagon-like peptide-1 receptor agonist treatment patterns among type 2 diabetes patients in six European countries. *Diabetes Ther* 2014; **5**: 499–520.
- National Institute for Health and Care Excellence. Type 2 diabetes—the management of type 2 diabetes (NICE clinical guideline 87, March 2014). London, 2014.
- Buse JB, Bergenstal RM, Glass LC et al. Use of twice-daily exenatide in basal insulin-treated patients with type 2 diabetes: a randomized, controlled trial. *Ann Intern Med* 2011; **154**: 103–112.
- Jensen MD, Ryan DH, Apovian CM et al. 2013 AHA/ACC/TOS guideline for the management of overweight and obesity in adults: a report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines and The Obesity Society. *Circulation* 2014; **129**: S102–S138.
- Tsigos C, Hainer V, Basdevant A et al. Management of obesity in adults: european clinical practice guidelines. *Obes Facts* 2008; **1**: 106–116.
- Pencek R, Blickensderfer A, Li Y, Brunell SC, Anderson PW. Exenatide twice daily: analysis of effectiveness and safety data stratified by age, sex, race, duration of diabetes, and body mass index. *Postgrad Med* 2012; **124**: 21–32.
- Rosenstock J, Shenouda SK, Bergenstal RM et al. Baseline factors associated with glycemic control and weight loss when exenatide twice daily is added to optimized insulin glargine in patients with type 2 diabetes. *Diabetes Care* 2012; **35**: 955–958.
- Chitnis AS, Ganz ML, Benjamin N, Langer J, Hammer M. Clinical effectiveness of liraglutide across body mass index in patients with type 2 diabetes in the United States: a retrospective cohort study. *Adv Ther* 2014; **31**: 986–999.
- Eto K, Naito Y, Seino Y. Evaluation of the efficacy and safety of lixisenatide add-on treatment to basal insulin therapy among T2DM patients with different body mass indices from GetGoal trials. *Diabetol Metab Syndr* 2015; **7**: 106.
- Rodbard HW, Cariou B, Zinman B et al. Comparison of insulin degludec with insulin glargine in insulin-naïve subjects with type 2 diabetes: a 2-year randomized, treat-to-target trial. *Diabet Med* 2013; **30**: 1298–1304.
- Wang C, Mamza J, Idris I. Biphasic vs basal bolus insulin regimen in type 2 diabetes: a systematic review and meta-analysis of randomized controlled trials. *Diabet Med* 2015; **32**: 585–594.
- Origin Trial Investigators. Predictors of nonsevere and severe hypoglycemia during glucose-lowering treatment with insulin glargine or standard drugs in the ORIGIN trial. *Diabetes Care* 2015; **38**: 22–28.